

# NEUROFISIODEL® EMULGEL

## Topical innovation in peripheral Neuropathies and Neuralgia

with



**OZOILE®**  
ERBAGIL PATENT  
Stable Ozonides

### MAIN COMPONENTS

Ozoile® (Stable Ozonides with Vitamin E acetate), Capsicum Annum Fruit Extract, Harpagophytum Procumbens Root Extract, Arnica Montana Flower Extract, Ribes Nigrum Leaf Extract, Arginine, Valine, Isoleucine, Leucine, Glutamine, Thioctic Acid, Panthenol.

### OZOILE®

Acts against the inflammatory and itchy status. Counteracts oxidative stress and it has a protective free radicals scavenger action on peripheral nerve fibres of the skin. It creates an unfavorable environment to the microbiota alterations. Acts against redness, itching, burning, swelling and degenerative processes in progress. It transfers oxygen to the tissue restoring the correct physiology.

### VITAMIN E ACETATE

Soothing effect, it contrasts membrane lipid peroxidation. Blocks free radicals.

### CAPSICUM ANNUM FRUIT EXTRACT

It acts against the alterations of somato-sensory areas and the peripheral sensitization

### DEVIL'S CLAW, ARNICA MONTANA FLOWER EXTRACT, RIBES NIGRUM LEAF EXTRACT

Enhances the effectiveness of Ozoile® against inflammation and pain.

### ARGININE, VALINE, ISOLEUCINE, LEUCINE

Improve muscular tone and act against the atrophy

### GLUTAMINE

Inhibiting action of the peripheral neurotransmission

### THIOCTIC ACID

It acts against degenerative processes of oxidative nature

### PANTHENOL

Protects and nourishes the skin and nerve fibres in association with Ozoile®.

### INDICATIONS

For topical use in the treatment of peripheral neuropathies and neuralgia of various etiologies. It acts on pain, itching and tingling sensation.

### HOW TO USE

Topical use. Apply on intact skin, keeping away from mucosa. Do not apply to damaged skin and mucosa. Apply the product directly on the affected area and massage until completely absorbed.

It is recommended to apply the product twice a day as recommended by physician. The treatment period differs according to the conditions to be treated and the rapidity of therapeutic response.

### WARNINGS

External use. Apply on intact skin, keeping away from mucosa. Do not apply on damaged skin and mucosa. Do not ingest. Keep out of the reach of children. Do not use it if the packaging appears to be damaged. Do not use the device beyond the expiration date. The expiration date refers to the intact and correctly stored product. In case of contact with eyes: rinse thoroughly for several minutes. Remove contact lenses if easy to do. Continue rinsing. Store in a cool, dry place away from heat sources and direct sunlight. Store preferably between 5 °C and 35 °C.

Wash your hands well before using the medical device. Carefully close the tube after use. After opening it is recommended to use the product within 6 months. Neurofisiodel® emulgel is well tolerated. It is contraindicated in case of ascertained hypersensitivity towards the components. There are no cases of sensitization or side effects if this phenomenon occurs please discontinue use and consult your physician.

### PACKAGE

40 ml - 1.4 fl.Oz. tube

GLUTEN FREE

NO COLOURS

NO ALLERGENS

 ERBAGIL s.r.l.  
Via L. Settebrini, 13  
82037 Telesio Terme (BN) Italy  
[www.erbagil.com](http://www.erbagil.com)



# NEUROFISIODOL® EMULGEL

## Lokalna inovacija pri perifernih nevropatijah in nevralgijah

Z



OZOILE®  
ERBAGIL PATENT  
Stabilni ozonidi

### GLAVNE SESTAVINE

Ozoile® (Stabilni Ozonidi z vitaminom E acetatom), Capsicum annum sadni ekstrakt, Harpagophytum Procumbens ekstrakt korenine, Arnica Montana cvetni ekstrakt, Ribes Nigrum ekstrakt listov, arginin, valin, izolevcin, levcin, glutamin, tioktična kislina, pantenol.

### OZOILE®

Deluje proti vnetno in proti srbenju. Preprečuje oksidativni stres in ima zaščitni učinek odstranjevanja prostih radikalov na periferna živčna vlakna kože. Ustvarja neugodno okolje za spremembe mikrobiote. Deluje proti nastajanju rdečice, srbenju, pekočem občutku, oteklinam in degenerativnim procesom. Prenaša kisik v tkivo in obnavlja pravilno fiziologijo.

### VITAMIN E ACETAT

Pomirjujoč učinek, nasprotuje membranski peroksidaciji lipidov. Blokira proste radikale.

### CAPSICUM ANNUUM SADNI EKSTRAKT

Deluje proti spremembam somatosenzoričnih področij in periferne senzibilizacije

### HUDIČEV KREMPELJ, ARNICA MONTANA, RIBEZ NIGRUM

Izboljšujejo učinkovitost mešanice Ozoile®, deluje proti vnetno in proti bolečinsko.

### ARGININ, VALIN, IZOLEVCIN, LEVCIN

Izboljša mišični tonus in deluje proti atrofiji

### GLUTAMIN

Inhibira delovanje periferne nevrotransmisije

### TIOKTIČNA KISLINA

Deluje proti degenerativnim in oksidacijskim procesom

### PANTENOL

V kombinaciji z Ozoile® ščiti in neguje kožo in živčna vlakna.

### INDIKACIJE

Za lokalno uporabo pri perifernih nevropatijah in nevralgijah različnih etiologij. Deluje na bolečino, srbenje in mravljinčenje.

### UPORABA

Lokalna uporaba. Nanesite na nepoškodovano kožo, pri čemer se izogibajte sluznice. Ne nanašajte na poškodovano kožo in sluznico. Izdelek nanesite neposredno na prizadeto področje in masirajte, dokler se popolnoma ne vpije. Priporočljivo je, da izdelek nanesete dvakrat na dan ali kot pripomočka zdravnik. Obdobje uporabe se razlikuje glede na hitrost terapevtskega odziva.

### OPOZORILA

Za zunanjou uporabo. Nanesite na nepoškodovano kožo, pri čemer se izogibajte sluznice. Ne nanašajte na poškodovano kožo in sluznico. Ne zaužijte. Hranite izven dosega otrok. Ne uporabljajte, če embalaža izgleda poškodovana. Pripomočka ne uporabljajte po preteku datuma veljavnosti. Rok uporabnosti se nanaša na nepoškodovan in pravilno shranjen izdelek. V primeru stika z očmi: temeljito izpirati nekaj minut. Odstranite kontaktne leče, če je to možno. Nadaljujte z izpiranjem. Shranujte na hladnem, suhem mestu, stran od virov topote in stran od neposredne sončne svetlobe. Shranujte pri temperaturi med 5 °C in 35 °C. Pred uporabo medicinskega pripomočka si dobro umijte roke. Po uporabi tubo previdno zaprite. Po odprtju je priporočljivo porabiti izdelek v roku 6 mesecev. Neurofisiadol® emulgel pacienti dobro prenašajo. Kontraindiciran je v primeru ugotovljene preobčutljivosti na sestavine. Primeri preobčutljivosti oz. stranski učinki niso znani, če pa se vseeno pojavijo, je priporočljivo takoj prenehati z uporabo in se posvetovati s svojim zdravnikom.

### PAKIRANJE

Tuba 40 ml

BREZ GLUTENA

BREZ BARVIL

BREZ ALERGENOV

Distributer:  
MagPharm,  
prodaja farmacevtskih izdelkov, d.o.o.  
Litostroška 44,  
1000 LUBLJANA

ERBAGIL s.r.l  
Via L. Settembrini, 13  
82037 Telesse Terme (BN) Italy  
[www.erbagil.com](http://www.erbagil.com)

